

Suicide Prevention in the U.S.

Each year, suicide accounts for more than 38,000 deaths in the United States.¹ In 2011, it was the 10th leading cause of death in the U.S.² The loss of someone to suicide resonates among family, friends, coworkers, and others in the community; it has been estimated that for each person who commits suicide, 5 to 10 other people are severely affected by the loss.³ Family and friends may experience a range of painful emotions, such as shock, anger, guilt, and depression.⁴ Suicide can occur across demographics, but certain groups are more at risk than the general population. Problems with intimate partner relationships and mental and physical health problems are factors that have been associated with the occurrence of suicide.⁵


Suicide is devastating, but there are resources and information available to help prevent it. **Learn more to be the one who makes a difference.**



Suicide claims the life of someone in the United States every 15 minutes.⁶

8.5
million

In 2009, an estimated 8.5 million adults aged 18 or older **reported having serious thoughts of suicide.**⁷



Women attempt suicide more often than men,⁸ yet, **men die more often** from suicide attempts. Men represent 79 percent of all suicides in the U.S.⁹

\$34.6
BILLION
per
YEAR

Suicide results in an estimated \$34.6 billion in combined medical and work loss costs.¹⁰

Who is at higher risk of suicide?

Teenagers and young adults

Suicide is the third leading cause of death for 15 to 24 year olds.¹¹

Military Service Members

On average, a member of the Armed Forces commits suicide every 36 hours.¹²

Veterans

Suicide is the cause of death of an estimated 18 U.S. veterans each day.¹³

American Indians/Alaskan Natives

Rates of suicide among American Indian/Alaskan Natives ages 15 to 34 are more than double the national average for that age group.¹⁴

Gay, lesbian and bisexual youth


Gay, lesbian, and bisexual adolescents in grades 7 through 12 are more than twice as likely to attempt suicide as heterosexuals within the same age range.¹⁵

Latina girls

Latina girls between the ages of 12 and 17 have the highest rate of suicide attempts among all adolescent ethnic groups.¹⁶

Had Serious Thoughts of Suicide in the Past Year Among People Aged 18 or Older, by State:

Percentages, Annual Averages Based on 2010 and 2011 NSDUHs



Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2010 (Revised March 2012) and 2011. <http://www.samhsa.gov/data/NSDUH/2k11State/NSDUHsaeMaps2011.htm>

A National Perspective

Suicide prevention in every state and community is important — even the loss of one life is too many. Though suicide can occur in any community, some regions of the U.S. are more affected than others. No matter where you live, there are steps that can help prevent suicides and make a positive change in your state and community. See below to find out what you can do.

What Communities Can Do

Families, Friends, Educators, Coworkers and Other Community Members

Learn and recognize the warning signs of suicide, including:¹⁷

- Talking about: wanting to die or kill oneself, being a burden to others, feeling hopeless or having no reason to live, or feeling trapped or in unbearable pain.
- Looking for a way to kill oneself, such as searching online or buying a gun.
- Behaving recklessly.
- Withdrawing or feeling isolated, displaying extreme mood swings, or increasing the use of alcohol or drugs.
- Showing rage or talking about seeking revenge, acting anxious or agitated, or sleeping too little or too much.

You can help someone who is considering suicide by taking one or more of the following steps:¹⁸

- Be willing to listen, and don't lecture or debate whether suicide is right or wrong. Allow the person in need to express his or her feelings without passing judgment or acting shocked.

- Be direct and get involved. Talk openly and matter-of-factly, showing interest and support for the person at risk.
- Don't be sworn to secrecy — seek support and professional help for the person at risk.
- Offer hope that alternatives are available.
- Get help from people or agencies that specialize in crisis intervention and suicide prevention.
 - Seek confidential help, available 24 hours a day, through the **National Suicide Prevention Lifeline: 1-800-273-TALK (8255)**, a toll-free service funded by SAMHSA.

Community Coalitions and Organizations

- Meet with local officials and ask them to support suicide prevention in the community and statewide.¹⁹
- Facilitate trainings on different topics, such as intervention skills, for key stakeholders and community members.²⁰
- Provide informational materials to schools or at local events.²¹

School Administrators

- Use SAMHSA's **Preventing Suicide: A Toolkit for High Schools**, available at <http://store.samhsa.gov/product/Preventing-Suicide-A-Toolkit-for-High-Schools/SMA12-4669>, for information on how to implement a suicide prevention program that responds to the needs and cultures of students.
- Help prevent suicide by establishing comprehensive crisis plans and school-based prevention and anti-bullying programs. See below for resources that can provide additional information. Learn more about national efforts in the **National Prevention Strategy**, available at <http://www.healthcare.gov/prevention/nphpphc/strategy/report.pdf> and the **2012 National Strategy for Suicide Prevention**, available at http://www.surgeongeneral.gov/library/reports/national-strategy-suicide-prevention/full_report-rev.pdf.

Parents and Caregivers

- Ask your child's friends and teachers if you have concerns about your child's behavior.
- Spend at least 15 minutes per day listening and talking to your children. Ask about their concerns, and express what you've heard back to them to make sure you understand each other.

Workplace²²

- Inform employees about the warning signs, risk factors, and prevention approaches for suicide.
- Create a work environment in which co-workers care for and support each other.
- Promote the development of interpersonal and social competency through training in stress management and coping skills to help individuals deal with problems.
- Expand awareness of suicide-prevention programs and mental health and substance use disorder assessment and treatment.

Helpful Resources

- **The National Suicide Prevention Lifeline** website, <http://www.suicidepreventionlifeline.org>, provides a downloadable wallet card with the Lifeline phone number and suicide warning signs in English and Spanish, as well as other materials for coping and caring for loved ones.
- The **StopBullying.gov** website, <http://www.stopbullying.gov>, provides information from various government agencies on what bullying is, what cyberbullying is, who is at risk, and how you can prevent and respond to bullying.
- **SAMHSA's Mental Health Services Locator**, available at <http://store.samhsa.gov/mhlocator>, can help people find mental health services and resources in their communities.
- For broader policy issues, the **National Action Alliance for Suicide Prevention**, available at <http://www.actionallianceforsuicideprevention.org>, is a public-private collaboration SAMHSA has developed to help promote suicide prevention.
- **The Trevor Project**, available at <http://www.thetrevorproject.org>, promotes acceptance of gay, lesbian, bisexual, and questioning teens and helps to prevent suicide among those youth. **The Trevor Helpline**, which can be reached at 1-866-488-7386, is a 24-hour toll-free suicide helpline for gay, lesbian, bisexual, and questioning youth.
- **The Suicide Prevention Resource Center**, available at <http://www.sprc.org>, provides access to the science and experience that can support efforts to develop programs, implement interventions, and promote policies to prevent suicide. Resources include information on school-based prevention programs, a best practices registry, state information and more.

Confidential help is available 24 hours a day through the National Suicide Prevention Lifeline: 1-800-273-TALK (8255), a toll-free service funded by SAMHSA.

You can also call SAMHSA's National Helpline, 1-800-662-HELP (4357) or 1-800-487-4889 (TDD), for 24-hour free and confidential information and treatment referrals in English and Spanish.

Sources

- ¹ Centers for Disease Control and Prevention (CDC). (2012). *Deaths: Preliminary Data for 2011*. (National Vital Statistics Report, Vol. 61, No. 6: October 10, 2012). Retrieved November 19, 2012, from http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_06.pdf
- ² Centers for Disease Control and Prevention (CDC). (2012). *Deaths: Preliminary Data for 2011*. (National Vital Statistics Report, Vol. 61, No. 6: October 10, 2012). Retrieved November 19, 2012, from http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_06.pdf
- ³ Substance Abuse and Mental Health Services Administration (SAMHSA). (2006). *CSAP's Prevention Pathways: Holding the Lifeline, a Guide to Suicide Prevention*. Retrieved December 20, 2011, from <https://www.pmrts.samhsa.gov/PrevCourses/>.
- ⁴ Centers for Disease Control and Prevention (CDC). (2010). *Suicide: Facts at a Glance*. Retrieved December 20, 2011, from http://www.cdc.gov/violenceprevention/pdf/Suicide_DataSheet-a.pdf.
- ⁵ Centers for Disease Control and Prevention (CDC). (2008). *Surveillance for Violent Deaths – National Violent Death Reporting System, 16 States, 2005*. (Center for Disease Control and Prevention, Surveillance Summaries, 57(SS03); 1-43, 45). Retrieved December 20, 2011, from <http://www.cdc.gov/mmwr/preview/mmwrhtml/ss5703a1.htm>
- ⁶ Centers for Disease Control and Prevention (CDC). (2010). *Web-based Injury Statistics Query and Reporting System (WISQARS) (Online)*. Available from www.cdc.gov/injury/wisqars/index.html. As cited in Centers for Disease Control and Prevention (CDC). (2010). *Suicide: Facts at a Glance*. Retrieved December 20, 2011, from http://www.cdc.gov/violenceprevention/pdf/Suicide_DataSheet-a.pdf
- ⁷ Substance Abuse and Mental Health Services Administration. (2012). *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*. NSDUH Series H-45, HHS Publication No. (SMA) 12-4725. Rockville, MD: Substance Abuse and Mental Health Services Administration.
- ⁸ Suicide Prevention Resource Center. *Suicide Prevention Basics: About Suicide*. Retrieved November 21, 2012, from <http://www.sprc.org/basics/about-suicide>
- ⁹⁻¹⁰ Centers for Disease Control and Prevention (CDC). (2010). *Suicide: Facts at a Glance*. Retrieved December 20, 2011, from http://www.cdc.gov/violenceprevention/pdf/Suicide_DataSheet-a.pdf. As cited in Centers for Disease Control and Prevention (CDC). (2010). *Web-based Injury Statistics Query and Reporting System (WISQARS) (Online)*. (2010). Retrieved December 20, 2011, from <http://www.cdc.gov/injury/wisqars/index.html>
- ¹¹ Centers for Disease Control and Prevention (CDC). (2007). *Web-based Injury Statistics Query and Reporting System (WISQARS) (Online)*. Retrieved November 20, 2012, from <http://www.cdc.gov/injury/wisqars/index.html>.
- ¹² Department of Defense Task Force on the Prevention of Suicide by Members of the Armed Forces. (2010). *Final report of the Department of Defense Task Force on the Prevention of Suicide by Members of the Armed Forces*. Retrieved December 20, 2011, from <http://www.health.mil/dhb/downloads/Suicide%20Prevention%20Task%20Force%20final%20report%208-23-10.pdf>. As cited in Substance Abuse and Mental Health Services Administration (SAMHSA). (2011). *Leading Change: A Plan for SAMHSA's Roles and Actions 2011-2014*. (HHS Publication No. SMA 11-4629). Rockville, MD: SAMHSA.
- ¹³ Department of Veterans Affairs (VA). (2010). *Remarks by Secretary Eric K. Shinseki*. (Office of Public and Intergovernmental Affairs, Suicide Prevention Conference, January 11, 2010). Washington, DC: VA. Retrieved December 20, 2011, from http://www.va.gov/opa/speeches/2010/10_0111hold.asp
- ¹⁴ Centers for Disease Control and Prevention (CDC). (2010). *Suicide: Facts at a Glance*. Retrieved December 20, 2011, from http://www.cdc.gov/violenceprevention/pdf/Suicide_DataSheet-a.pdf. As cited in Centers for Disease Control and Prevention (CDC). (2010). *Web-based Injury Statistics Query and Reporting System (WISQARS) (Online)*. (2010). Retrieved December 20, 2011, from <http://www.cdc.gov/injury/wisqars/index.html>
- ¹⁵ Russell ST, Joyner K. (2001). Adolescent sexual orientation and suicide risk: Evidence from a national study. *American Journal of Public Health*, 91:1276–1281. As cited by Centers for Disease Control and Prevention (CDC). (2011). *Lesbian, Gay, Bisexual and Transgender Health*. Retrieved December 20, 2011, from <http://www.cdc.gov/lgbthealth/youth.htm>
- ¹⁶ Centers for Disease Control and Prevention. *Youth Risk Behavior Surveillance — United States, 2007*. Surveillance Summaries, June 6. MMWR 2008; 57(No. SS-4). As cited in Centers for Disease Control and Prevention. (2009). *Suicide: Facts at a Glance*. Retrieved November 20, 2012, from <http://www.cdc.gov/violenceprevention/pdf/suicide-datasheet-a.pdf>
- ¹⁷ Substance Abuse and Mental Health Services Administration (SAMHSA). (2011). *National Suicide Prevention Lifeline. Learn the Warning Signs*. (HHS Publication No. SVP-0126). Retrieved December 20, 2011, from http://www.suicidepreventionlifeline.org/App_Files/Media/PDF/NSPL_WalletCard.pdf
- ¹⁸ Substance Abuse and Mental Health Services Administration (SAMHSA). *National Suicide Prevention Lifeline. Get Help for Someone Else*. Retrieved December 20, 2011, from <http://www.suicidepreventionlifeline.org/GetHelp/Someone>
- ¹⁹ The Massachusetts Coalition for Suicide Prevention. *Advocacy*. Retrieved December 20, 2011, from <http://www.masspreventsuicide.org/advocacy>
- ²⁰ Suicide Prevention Coalition of Colorado. *Calendar of Events*. Retrieved December 20, 2011, from <http://www.suicidepreventioncolorado.org/calendar-of-events.html>
- ²¹ Indiana Suicide Prevention Coalition. Retrieved December 20, 2011, from <http://www.indianasuicidepreventioncoalition.org>
- ²² Great-West Life Centre for Mental Health in the Workplace. (2011). *Workplace Strategies for Mental Health – Suicide Intervention*. Retrieved on December 11, 2012, from <http://www.gwlcetreformentalhealth.com/display.asp?l1=7&l2=101&l3=116&d=116#1>

